



Vaccine Fact Sheet

HPV9

Brand Name and Manufacturer	Gardasil®9 Merck
Protects Against	Human papillomavirus types 6, 11, 16, 18, 31, 33, 45, 52, and 58 which can cause cervical, vulvar; vaginal, and anal cancers, as well as genital warts.
Routine Schedule	Two (2) doses for persons initiating vaccination at age 9 through 14 years, except immunocompromised persons Three (3) doses for persons initiating vaccination at age 15 through 26 years, and immunocompromised persons initiating vaccination at age 9 through 26 years
Minimum Intervals	Two-dose series: · 5 months between doses Three-dose series: · 4 weeks between doses 1 and 2 · 12 weeks between doses 2 and 3 · 5 months between doses 1 and 3
Approved for use in	Females and males ages 9 through 26 years
Administration	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single-dose 0.5mL vials
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) PROTECT VACCINE FROM LIGHT DO NOT FREEZE
Full ACIP Recommendations	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/hpv.html
VFC Letter	http://eziz.org/assets/docs/VFC_Letters/Feb162017HPV2dose.pdf
Billing Codes	CHDP code: 93 CPT code for vaccine: 90651 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90651 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * http://www.aafp.org/practice-management/payment/coding/admin.html
Comments	<ul style="list-style-type: none">• The 9-valent HPV vaccine may be used to complete a series previously started with another HPV vaccine product.• The bivalent and quadrivalent HPV vaccines are no longer distributed in the U.S.• ACIP recommends routine vaccination with HPV vaccine at age 11 or 12 years. The vaccination series may be initiated as early as age 9 years.• The minimum intervals for HPV vaccination should only be used retrospectively to validate doses. Use the recommended intervals when scheduling patients for their next HPV vaccine dose.• Syncope (fainting) can occur after vaccination, vaccine recipients should be observed for 15 minutes after vaccination. <p>Regardless of licensure, vaccines available through the VFC Program can only be used for VFC-eligible children through 18 years of age.</p>