



Vaccine Fact Sheet

PCV 13

Brand Name and Manufacturer	Prevnar 13® Wyeth/Pfizer
Protects Against	13 types of pneumococcal bacteria which can cause meningitis, blood infections and pneumonia
Routine Schedule	Four (4) dose series: 2, 4, 6 and 12-15 months
Minimum Intervals	For children <12 months, 4 week minimum interval between doses For children >12 months, 8 weeks between doses
Approved for use in	Children aged 6 weeks through 5 years
Administration	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single-dose 0.5mL pre-filled disposable syringes
Storage	Refrigerate between 35°F and 46°F (2°C to 8°C) DO NOT FREEZE
Full ACIP Recommendations	http://www.cdc.gov/vaccines/recs/provisional/downloads/pcv13-mar-2010-508.pdf
VFC Letter	http://www.eziz.org/PDF/13ValentPneumococcalConjugateVac.pdf
Billing Codes	CHDP code: 88 CPT code for vaccine: 90670 CPT code for administration: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90670-SL ICD-9-CM code: v03.82
Comments	<ul style="list-style-type: none">• Licensed in 2010• Recommended for children aged 60 months through 71 months with increased risk of pneumococcal disease or its complications• Recommended for children aged 6 years through 18 years with sickle cell disease, HIV infection or other immunocompromising conditions, cochlear implants or CSF leaks may receive a dose of PCV13.• A single supplemental dose is recommended for all children aged 14 through 59 months who received an age-appropriate series of Prevnar 7.