



Vaccine Fact Sheet

PCV13

Brand Name and Manufacturer	Pevnar 13® Wyeth/Pfizer
Protects Against	13 types of pneumococcal bacteria which can cause meningitis, blood infections and pneumonia
Routine Schedule	Four (4) dose series: 2, 4, 6 and 12-15 months
Minimum Intervals	For children <12 months, 4 week minimum interval between doses For children >12 months, 8 weeks between doses
Approved for use in	Children aged 6 weeks through 5 years
Administration	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single-dose 0.5mL pre-filled disposable syringes
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE
Full ACIP Recommendations	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/pneumo.html
VFC Letter	http://www.eziz.org/PDF/13ValentPneumococcalConjugateVac.pdf
Billing Codes*	CHDP code: 88 CPT code for vaccine: 90670 CPT code for administration: 90640 Medi-Cal Fee-For-Service (FFS) administration: 90670 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 *http://www.aafp.org/practice-management/payment/coding/admin.html
Comments	<ul style="list-style-type: none">• Licensed in 2010.• DO NOT administer PCV13 and PPSV23 at the same visit.• For children ages 2-18 years, refer to Pneumococcal Vaccine Timing—For Children, IMM-1159.• Routinely recommended for children ages 2-59 months.• Recommended for children ages 60 months through 71 months with increased risk of pneumococcal disease or its complications.• Recommended for children ages 6 years through 18 years with sickle cell disease, HIV infection or other immunocompromising conditions, cochlear implants or CSF leaks.• For adults ages 19 and older, refer to Pneumococcal Vaccine Timing—For Adults, IMM-1152.