



Vaccine Fact Sheet

Hib-HepB Combination Vaccine

Protects Against *Haemophilus influenzae* type b (Hib) and Hepatitis B

Brand Name and Manufacturer COMVAX®
Merck

Routine Schedule Three (3) dose series: 2, 4, and 12-15 months

Minimum Intervals 4 week minimum interval between dose 1 and 2
8 week minimum interval between dose 2 and 3
(the third dose should not be given before 12 months of age)

Approved for use in Children aged 6 weeks to 15 months

Administration Intramuscular (IM) injection

Packaging Vaccine is packaged as 10 single-dose 0.5mL vials

Storage Refrigerate between 35°F and 46°F (2°C to 8°C)

DO NOT FREEZE

Full ACIP Recommendations Hib:
<http://www.cdc.gov/mmwr/preview/mmwrhtml/00041736.htm>
Hep B:
http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5416a1.htm?s_cid=rr5416a1_e

VFC Letter http://eziz.org/assets/docs/vfcletter_2012_03_RecommendedACIPschedules.pdf

Billing Codes CHDP code: 56
CPT code for vaccine: 90748
CPT code for administration: 90460, 90461
Medi-Cal Fee-For-Service (FFS) administration: 90748-SL
ICD-9-CM code: v06.8

Comments

- Licensed in 1996
- Do not administer to infants younger than 6 weeks of age.
- Infants born to HBsAG-positive mothers should receive Hepatitis B Immune Globulin and Hepatitis B vaccine within 12 hours of birth and should complete the Hepatitis B vaccination series according to schedule.