



Vaccine Fact Sheet

IPV

Protects Against Poliovirus

Brand Name and Manufacturer IPOL®
Sanofi Pasteur

Routine Schedule Four dose (4) series: 2, 4, and 6-18 months, and 4-6 years

Minimum Intervals 4 week minimum interval between dose 1 and 2
4 week minimum interval between dose 2 and 3
6 month minimum interval between dose 3 and 4

Approved for use in Persons aged 6 weeks and older

Administration Intramuscular (IM) injection or Subcutaneous (SC) injection

Packaging Vaccine is packaged as a 10-dose multi-dose vial

Storage Refrigerate between 35°F and 46°F (2°C to 8°C)
DO NOT FREEZE

Full ACIP Recommendations <http://www.cdc.gov/vaccines/pubs/pinkbook/polio.html>

VFC Letter <http://eziz.org/assets/docs/IMM-1061.pdf>

Billing Codes CHDP code: 39
CPT code for vaccine: 90713
CPT code for administration: 90460
Medi-Cal Fee-For-Service (FFS) administration: 90713-SL
ICD-9-CM code: v04.0

- Comments**
- Licensed in 1990
 - In the first 6 months of life, the minimum age and minimum intervals are only recommended if the infant is at risk of imminent exposure to the disease (i.e., travel to a polio-endemic region or during an outbreak).
 - A fourth dose is not necessary if the third dose was given at 4 years of age or older and at least 6 months after the second dose.
 - IPV is not routinely recommended for U.S. residents aged 18 years or older.