



Vaccine Fact Sheet

MMR

Protects Against Measles, Mumps, and Rubella

Brand Name and Manufacturer M-M-R II®
Merck

Routine Schedule Two (2) dose series: First dose at 12-15 months and second dose at 4-6 years

Minimum Intervals 4 week minimum interval between dose 1 and 2

Approved for use in Persons aged 12 months and older

Administration Subcutaneous (SC) injection

Packaging Vaccine is packaged as 10 single-dose vials of lyophilized vaccine and a separate box of 10 single-dose 0.7mL vials of sterile diluent

Storage Store between -58°F and +46°F (-50°C to +8°C)
RECOMMEND STORAGE IN FREEZER
PROTECT VACCINE FROM LIGHT

Full ACIP Recommendations <http://www.cdc.gov/mmwr/preview/mmwrhtml/00053391.htm>

VFC Letter Not available on EZIZ

Billing Codes CHDP code: 33
CPT code for vaccine: 90707
CPT code for administration: 90460
Medi-Cal Fee-For-Service (FFS) administration: 90707 with modifiers –SK (high-risk) and –SL (VFC)
ICD-10-CM code (encounter for immunization): Z23
[*http://www.aafp.org/practice-management/payment/coding/admin.html](http://www.aafp.org/practice-management/payment/coding/admin.html)

- Comments**
- Licensed in 1971
 - Give a second dose to all older children and teens with history of only 1 dose.
 - Give MMR to infants aged 6 through 11 months who are traveling internationally. This dose does not count toward the 2-dose routine series and these children should be revaccinated at ages 12-15 months (at least 4 weeks after previous dose) with the second dose at ages 4-6 years.
 - MMR vaccine must be used within 8 hours of reconstitution.
 - Use only the Merck sterile diluent to reconstitute lyophilized MMR vaccine.
 - Store sterile diluent separately at room temperature (68°F to 77°F) or in the refrigerator.
 - If MMR, varicella, and/or LAIV are not administered at the same visit, they should be separated by at least 28 days.
 - Persons who are immunocompromised due to certain conditions or treatments or who are pregnant should not be vaccinated with MMR vaccine.