



Vaccine Fact Sheet

DTaP

Protects Against	Diphtheria, tetanus, and pertussis	Diphtheria, tetanus, and pertussis
Brand Name and Manufacturer	Infanrix® GlaxoSmithKline (GSK)	Daptacel® Sanofi Pasteur
Routine Schedule	Three (3) dose series: 2, 4, and 6 months Two (2) booster doses: 15-18 months and 4-6 years	Three (3) dose series: 2, 4, and 6 months Two (2) booster doses: 15-18 months and 4-6 years
Minimum Intervals	4 week minimum interval between dose 1 and 2 4 week minimum interval between dose 2 and 3 6 months minimum interval between dose 3 and 4 6 months minimum interval between dose 4 and 5	4 week minimum interval between dose 1 and 2 4 week minimum interval between dose 2 and 3 6 months minimum interval between dose 3 and 4 6 months minimum interval between dose 4 and 5
Approved for use in	Children aged 6 weeks through 6 years	Children aged 6 weeks through 6 years
Administration	Intramuscular (IM) injection	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single-dose 0.5mL vials or 10 single-dose 0.5mL pre-filled disposable syringes without needles.	Vaccine is packaged as 10 single-dose 0.5mL vials
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE
Full ACIP Recommendations	www.cdc.gov/mmwr/preview/mmwrhtml/00048610.htm	www.cdc.gov/mmwr/preview/mmwrhtml/00048610.htm www.cdc.gov/mmwr/preview/mmwrhtml/mm5126a5.htm
VFC Letter	Not available on EZIZ	Not available on EZIZ
Billing Codes	CHDP code: 45 CPT code for vaccine: 90700 CPT code for administration*: 90460, 90461 Medi-Cal Fee-For-Service (FFS) administration: 90644 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 *http://www.aafp.org/practice-management/payment/coding/admin.html	CCHDP code: 45 CPT code for vaccine: 90700 CPT code for administration*: 90460, 90461 Medi-Cal Fee-For-Service (FFS) administration: 90644 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 *http://www.aafp.org/practice-management/payment/coding/admin.html
Comments	<ul style="list-style-type: none"> Licensed in 1997 The fourth dose of the primary series may be given as early as 12 months of age provided at least 6 months have elapsed since the third dose If the fourth dose is given after the fourth birthday, the fifth dose is not necessary Do not give DTaP to children age 7 years or older; use Tdap If possible, use the same DTaP product for all doses 	<ul style="list-style-type: none"> Licensed in 2002 The fourth dose of the primary series may be given as early as 12 months of age provided at least 6 months have elapsed since the third dose If the fourth dose is given after the fourth birthday, the fifth dose is not necessary Do not give DTaP to children age 7 years or older; use Tdap If possible, use the same DTaP product for all doses

