## **Vaccine Fact Sheet: MMRV**



Topic	ProQuad®
Manufacturer	Merck <u>Detailed Prescribing Information</u> (Merck.com/product/usa/pi_circulars/p/proquad/proquad_pi_4171.pdf)
Protects Against	Measles, Mumps, Rubella and Varicella
Routine Schedule	Two (2) dose series: First dose at 12-15 months and second dose at 4-6 years.
Minimum Intervals	At least 1 month between a dose of a measles-containing vaccine, i.e. MMR, and a dose of ProQuad <sup>®</sup> . At least 3 months should elapse between a dose of varicella-containing vaccine and a dose of ProQuad <sup>®</sup> .
Approved Ages	Persons aged 12 months through 12 years of age
Administration	Subcutaneous (SC) or Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single-dose vials of lyophilized vaccine (Package A) and a separate package of 10 vials of sterile water diluent (Package B)
Storage	Store frozen between -58°F and +5°F (-50°C to -15°C).  Store in freezer  Protect vaccine from light
Full ACIP Recommendations	MMR Recommendations of the Advisory Committee on Immunization Practices (ACIP) (Cdc.gov/mmwr/pdf/rr/rr5903.pdf)
VFC Letter	VFC Clinical Letter (Eziz.org/assets/docs/vfcletter_2012_11_CombinedMMRV.pdf)
Billing Codes	CHDP code: 74 CPT code for vaccine: 90710 CPT code for administration with counseling under 8 years of age: 90460 or 90461 CPT code for administration without counseling under 8 years of age: 90471 or 90472 Medi-Cal Fee-For-Service (FFS) administration: 90707 with modifiers –SK (highrisk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 Child Health and Disability Prevention (CHDP) Program Code Conversion (Files.medical.ca.gov/pubsdoco/chdp/articles/25768.02_Cd_Conv_Table.pdf)
Comments	Licensed in 2005