


Administering Vaccines: Dose, Route, Site, and Needle Size

Vaccines	Dose	Route
Diphtheria, Tetanus, Pertussis (DTaP, DT, Tdap, Td)	0.5 mL	IM
<i>Haemophilus influenzae type b (Hib)</i>	0.5 mL	IM
Hepatitis A (HepA)	≤18 yrs: 0.5 mL ≥19 yrs: 1.0 mL	IM
Hepatitis B (HepB)	≤19 yrs: 0.5 mL* ≥20 yrs: 1.0 mL	IM
*Persons 11–15 yrs may be given Recombivax HB® (Merck) 1.0 mL adult formulation on a 2-dose schedule.		
Human papillomavirus (HPV)	0.5 mL	IM
Influenza, live attenuated (LAIV)	0.2 mL	Intranasal spray
Influenza, trivalent inactivated (TIV)	6–35 mos: 0.25 mL ≥3 yrs: 0.5 mL	IM
Measles, mumps, rubella (MMR)	0.5 mL	SC
Meningococcal, conjugated (MCV4)	0.5 mL	IM
Meningococcal, polysaccharide (MPSV4)	0.5 mL	SC
Pneumococcal conjugate (PCV)	0.5 mL	IM
Pneumococcal polysaccharide (PPV)	0.5 mL	IM or SC
Polio, inactivated (IPV)	0.5 mL	IM or SC
Rotavirus (RV)	2.0 mL	Oral
Varicella (Var)	0.5 mL	SC
Zoster (Zos)	0.65 mL	SC
Combination Vaccines		
DTaP+HepB+IPV (Pediarix™) DTaP+Hib (Trihibit™) Hib+HepB (Comvax™)	0.5 mL	IM
MMR+Var (ProQuad®)	≤12 yrs: 0.5 mL	SC
HepA+HepB (Twinrix®)	≥18 yrs: 1.0 mL	IM

Injection Site and Needle Size		
Subcutaneous (SC) injection		
Use a 23–25 gauge needle. Choose the injection site that is appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Infants (1–12 mos)	5/8"	Fatty tissue over anterolateral thigh muscle
Children (≥12 mos), adolescents, and adults	5/8"	Fatty tissue over anterolateral thigh muscle or fatty tissue over triceps
Intramuscular (IM) injection		
Use a 22–25 gauge needle. Choose the injection site and needle length appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Newborns (1 st 28 days)	5/8"*	Anterolateral thigh muscle
Infants (1–12 mos)	1"	Anterolateral thigh muscle
Toddlers (1–2 yrs)	1"–1¼" 5/8"**–1"	Anterolateral thigh muscle or deltoid muscle of arm
Children & teens 3–18 yrs	5/8"**–1" 1"–1¼"	Deltoid muscle of arm or anterolateral thigh muscle
Adults ≥ age 19 yrs		
Male or female less than 130 lbs	5/8"**–1"	Deltoid muscle of arm
Female 130–200 lbs Male 130–260 lbs	1"–1½"	Deltoid muscle of arm
Female 200+ lbs Male 260+ lbs	1½"	Deltoid muscle of arm

*If skin is stretched tight and subcutaneous tissue is not bunched.



Please note: Always refer to the package insert included with each biologic for complete vaccine administration information. CDC's Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well.