









## Routine Schedule

| Age*                 | Vaccine                    | Primary Doses  | Booster Dose |
|----------------------|----------------------------|--|--------------|
| 6 months–<br>4 years | Pfizer–<br>Infant/Toddler  |  |              |
| 6 months–<br>5 years | Moderna–<br>Infant/Toddler |   |              |
| 5–11 years           | Pfizer–<br>Pediatric       |  |              |
| 6–11 years           | Moderna–<br>Pediatric      |   |              |

## Schedule if Moderately or Severely Immunocompromised

| Age*                 | Vaccine                    | Primary Doses  | Booster Dose |
|----------------------|----------------------------|--|--------------|
| 6 months–<br>4 years | Pfizer–<br>Infant/Toddler  |  |              |
| 6 months–<br>5 years | Moderna–<br>Infant/Toddler |  |              |
| 5–11 years           | Pfizer–<br>Pediatric       |  |              |
| 6–11 years           | Moderna–<br>Pediatric      |  |              |

\* See schedules for children in transition from a younger to older age group: [Pfizer](#) | [Moderna](#).

^ An 8-week interval may be preferable for some people, especially for males 12-39 years.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

# COVID-19 Vaccine Timing–Adolescent/Adult

## Routine Schedule

| Age   | Vaccine                                    | Primary Doses  | Booster Doses  |
|-------|--|--|--|
| 12-17 | Moderna–<br>Adol/Adult                     | 1st Dose → 4-8 weeks <sup>^</sup> → 2nd Dose               |  |
| 12+   | Pfizer/<br>Adol/Adult                      | 1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose → ≥5 months → | <div style="border: 1px solid black; padding: 10px; background-color: #e0f2e0;"> <p><b>1st Booster</b></p> <p><b>Ages 12-17:</b><br/>Pfizer</p> <p><b>18+:</b><br/>Moderna/ Pfizer<br/>(mRNA preferred)<br/>or J&amp;J*</p> </div> <p>→ ≥4 months →</p> <div style="border: 1px solid black; padding: 10px; background-color: #e0f2e0;"> <p><b>2nd Booster</b></p> <p><b>Ages 50+:</b><br/>Moderna/Pfizer</p> <p><b>18-49:</b><br/>Not currently recommended.</p> <p>(If received J&amp;J for primary and 1st booster, may consider receiving a 2nd booster of mRNA vaccine.)</p> </div> |
| 18+   | Moderna–<br>Adol/Adult                     | 1st Dose → 4-8 weeks <sup>^</sup> → 2nd Dose → ≥5 months → |  |
| 18+   | Janssen (J&J)<br>Pfizer/Moderna preferred* | 1st Dose → ≥2 months →                                     |  |
| 18+   | Novavax                                    | 1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose               |  |







<sup>^</sup> An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

\* Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

# COVID-19 Vaccine Timing—Adolescent/Adult

## Schedule if Moderately or Severely Immunocompromised

| Age   | Vaccine                                       | Primary Doses   | Booster Doses   |
|-------|---|---|---|
| 12-17 | Moderna—<br>Adol/Adult                        |   |   |
| 12+   | Pfizer/<br>Adol/Adult                         |   | <div style="display: flex; align-items: center; justify-content: center;"> <div style="border: 1px solid #add8e6; border-radius: 15px; padding: 10px; text-align: center;"> <p><b>1st Booster</b></p> <p><b>Ages 12-17:</b><br/>Pfizer</p> <p><b>18+:</b><br/>Pfizer/<br/>Moderna<br/>(mRNA<br/>preferred)<br/>or J&amp;J*</p> </div> <div style="margin: 0 10px;">  </div> <div style="border: 1px solid #add8e6; border-radius: 15px; padding: 10px; text-align: center;"> <p><b>2nd Booster</b></p> <p><b>Ages 12-17:</b><br/>Pfizer</p> <p><b>18+:</b><br/>Moderna/<br/>Pfizer</p> </div> </div> |
| 18+   | Moderna—<br>Adol/Adult                        |   |   |
| 18+   | Janssen (J&J)<br>Pfizer/Moderna<br>preferred* |  |   |
| 18+   | Novavax                                       |  |   |

\* Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.