



# Vaccine Fact Sheet

## Rotavirus

<b>Brand Name and Manufacturer</b>	Rotarix® GlaxoSmithKline (GSK)	RotaTeq® Merck
<b>Protects Against</b>	Rotavirus	Rotavirus
<b>Routine Schedule</b>	Two (2) dose series, starting at 6-8 weeks of age	Three (3) dose series, starting at age 6-12 weeks of age
<b>Minimum Intervals</b>	4 weeks minimum interval between dose 1 and 2	4-10 weeks between dose 1 and 2; 4-10 weeks between dose 2 and 3
<b>Approved for use in</b>	Children aged 6 weeks to 24 weeks.	Children aged 6 weeks to 32 weeks. Do not administer to children older than 32 weeks.
<b>Administration</b>	Oral	Oral
<b>Packaging</b>	Vaccine is packaged as 10 single-dose vials of lyophilized vaccine, and 10 prefilled oral applicators of liquid (1 mL) with a plunger stopper and a transfer adapter.	Vaccine is packaged as 10 single-dose, ready-to-use 2mL pouches
<b>Storage</b>	Refrigerate between 36°F and 46°F (2°C to 8°C) <b>DO NOT FREEZE</b>	Refrigerate between 36°F and 46°F (2°C to 8°C) <b>DO NOT FREEZE</b>
<b>Full ACIP Recommendations</b>	<a href="https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html">https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html</a>	<a href="https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html">https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html</a>
<b>VFC Letter</b>	<a href="http://www.eziz.org/PDF/RotarixLive%20AttenuatedRotavirusVaccinedoc2Signed.pdf">http://www.eziz.org/PDF/RotarixLive%20AttenuatedRotavirusVaccinedoc2Signed.pdf</a>	<a href="http://www.cdph.ca.gov/programs/immunize/Documents/IZ%20Final%20Rotavirus%20VFC.pdf">http://www.cdph.ca.gov/programs/immunize/Documents/IZ%20Final%20Rotavirus%20VFC.pdf</a>
<b>Billing Codes</b>	CHDP code: 81 CPT code for vaccine: 90681 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90681 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * <a href="http://www.aafp.org/practice-management/payment/coding/admin.html">http://www.aafp.org/practice-management/payment/coding/admin.html</a>	CHDP code: 75 CPT code for vaccine: 90681 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90681 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * <a href="http://www.aafp.org/practice-management/payment/coding/admin.html">http://www.aafp.org/practice-management/payment/coding/admin.html</a>
<b>Comments</b>	<ul style="list-style-type: none"> <li>• Licensed in 2008</li> <li>• Vaccine must be reconstituted.</li> <li>• Do not begin series in infants older than 14 weeks, 6 days of age. Infants with severe latex allergies should not receive Rotarix.</li> <li>• Store the diluent for Rotarix® with the vaccine component.</li> <li>• Infants diagnosed with severe combined immunodeficiency disease (SCFD) or a history of intussusception should not receive Rotarix.</li> <li>• The lot number to be recorded should be taken from the label on the package/box.</li> </ul>	<ul style="list-style-type: none"> <li>• Licensed in 2006</li> <li>• Do not begin series in infants older than 14 weeks, 6 days of age. Infants with severe latex allergies should not receive RotaTeq.</li> <li>• Infants diagnosed with severe combined immunodeficiency disease (SCFD) or a history of intussusception should not receive RotaTeq.</li> </ul>