



Vaccine Fact Sheet

Rotavirus

Brand Name and Manufacturer	Rotarix® GlaxoSmithKline (GSK)	RotaTeq® Merck
Protects Against	Rotavirus	Rotavirus
Routine Schedule	Two (2) dose series, starting at 6-8 weeks of age	Three (3) dose series, starting at age 6-12 weeks of age
Minimum Intervals	4 weeks minimum interval between dose 1 and 2	4-10 weeks between dose 1 and 2; 4-10 weeks between dose 2 and 3
Approved for use in	Infants 6 weeks to 24 weeks of age. https://www.fda.gov/vaccines-blood-biologics/vaccines/rotarix	Infants 6 weeks to 32 weeks of age. https://www.fda.gov/vaccines-blood-biologics/vaccines/rota-teq
Administration	Oral	Oral
Packaging	Vaccine is packaged as 10 single-dose vials of lyophilized vaccine, and 10 prefilled oral applicators of liquid (1 mL) with a plunger stopper and a transfer adapter.	Vaccine is packaged as 10 single-dose, ready-to-use 2mL pouches
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE
Full ACIP Recommendations	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html Minimum age of first dose: 6 weeks and 0 days Maximum age of first dose: 14 weeks and 6 days Maximum age of last dose: 8 months and 0 days	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/rotavirus.html Minimum age of first dose: 6 weeks and 0 days Maximum age of first dose: 14 weeks and 6 days Maximum age of last dose: 8 months and 0 days
VFC Letter	http://www.eziz.org/PDF/RotarixLive%20AttenuatedRotavirusVaccinedoc2Signed.pdf	http://www.cdph.ca.gov/programs/immunize/Documents/IZ%20Final%20Rotavirus%20VFC.pdf
Billing Codes	CHDP code: 81 CPT code for vaccine: 90681 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90681 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * http://www.aafp.org/practice-management/payment/coding/admin.html	CHDP code: 75 CPT code for vaccine: 90681 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90681 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * http://www.aafp.org/practice-management/payment/coding/admin.html
Comments	<ul style="list-style-type: none"> Licensed in 2008 Vaccine must be reconstituted. Infants with severe latex allergies should not receive Rotarix. Store the diluent for Rotarix® with the vaccine component. Infants diagnosed with severe combined immunodeficiency disease (SCFD) or a history of intussusception should not receive Rotarix. The lot number to be recorded should be taken from the label on the package/box. 	<ul style="list-style-type: none"> Licensed in 2006 Infants with severe latex allergies should not receive RotaTeq. Do not begin series in infants older than 14 weeks, 6 days of age. Infants with severe latex allergies should not receive RotaTeq. Infants diagnosed with severe combined immunodeficiency disease (SCFD) or a history of intussusception should not receive RotaTeq.