



Vaccine Fact Sheet

MMRV

Protects Against Measles, Mumps, Rubella and Varicella

Brand Name and Manufacturer ProQuad® (MMRV)
Merck

Routine Schedule MMR and varicella vaccines at 12-15 months*; second dose MMRV at 4-6 years.

Minimum Intervals At least 1 month between a dose of a measles-containing vaccine, i.e. MMR, and a dose of ProQuad®. At least 3 months should elapse between a dose of varicella-containing vaccine and a dose of ProQuad®.

Approved for use in Persons aged 12 months through 12 years of age

Administration Subcutaneous (SC) injection

Packaging Vaccine is packaged as 10 single-dose vials of lyophilized vaccine (Package A) and a separate package of 10 vials of sterile water diluent (Package B)

Storage Store frozen between -58°F and +5°F (-50°C to -15°C).

STORE IN FREEZER

PROTECT VACCINE FROM LIGHT

Full ACIP Recommendations <http://www.cdc.gov/mmwr/pdf/rr/rr5903.pdf>

VFC Letter http://eziz.org/assets/docs/vfcletter_2012_11_CombinedMMRV.pdf

Billing Codes CHDP code: 74
CPT code for vaccine: 90710
CPT code for administration with counseling under 8 years of age: 90460 or 90461
CPT code for administration without counseling under 8 years of age: 90471 or 90472
Medi-Cal Fee-For-Service (FFS) administration: 90707 with modifiers –SK (high-risk) and –SL (VFC)
ICD-10-CM code (encounter for immunization): Z23
<http://www.aafp.org/practice-management/payment/coding/admin.html>

Comments

- Use only the Merck sterile diluent to reconstitute lyophilized vaccine.
- Use only the Merck sterile diluent to reconstitute lyophilized vaccine.
- Store sterile diluent separately at room temperature (68°F to 77°F) or in the refrigerator.
- Varicella vaccine must be used within 30 minutes of reconstitution.

* Unless parent or caregiver expresses a preference for MMRV as the first dose, administer MMR and varicella. For first doses given at 48 months or older, MMRV is generally preferred. See ACIP recommendations for details.