



Vaccine Fact Sheet

Td

Protects Against	Tetanus and diphtheria	Tetanus and diphtheria
Brand Name and Manufacturer	Tenivac® Sanofi Pasteur	Td Vaccine Mass Biologics/Grifols
Routine Schedule	Three (3) dose series	Three (3) dose series
Minimum Intervals	8 week minimum between dose 1 and 2 6 month minimum between dose 2 and 3	4 week minimum between dose 1 and 2 6 month minimum between dose 2 and 3
Approved for use in	Persons aged 7 years and older	Persons aged 7 years and older
Administration	Intramuscular (IM) injection	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 10 single dose vials or syringes.	Vaccine is packaged as 10 single dose vials
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE	Refrigerate between 36°F and 46°F (2°C to 8°C) DO NOT FREEZE
Full ACIP Recommendations	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/dtap.html	https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/dtap.html
VFC Letter	Td-specific letter not available on EZIZ	http://eziz.org/assets/docs/VFC_Letters/VFCletter-TdGrifols-Jan2017%20(2).pdf
Billing Codes	CHDPVFC: 58 CHDP Adult Purchased: 68 CPT code for vaccine: 90714 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90714 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * http://www.aafp.org/practice-management/payment/coding/admin.html	CHDPVFC: 58 CHDP Adult Purchased: 68 CPT code for vaccine: 90714 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90714 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 * http://www.aafp.org/practice-management/payment/coding/admin.html
Comments	<ul style="list-style-type: none"> • Several vaccines are used to prevent tetanus among children, adolescents, and adults including DTaP, Tdap, DT, and Td. • Tdap can be administered regardless of interval since the previous Td dose. • Do not administer Td vaccines to infants or children younger than 7 years of age. • Tdap can be used for the decennial Td booster. 	<ul style="list-style-type: none"> • Several vaccines are used to prevent tetanus among children, adolescents, and adults, including DTaP, Tdap, DT, and Td. • Tdap can be administered regardless of interval since the previous Td dose. • Do not administer Td vaccines to infants or children younger than 7 years of age. • Tdap can be used for the decennial Td booster.