

# Vaccine Fact Sheet: Td

Topic	TENIVAC®
Manufacturer	Sanofi Pasteur <a href="#">Prescribing Info (FDA)</a>
Protects Against	Tetanus & diphtheria
Approved Ages	7 years & older
Funding Source	<b>VFC</b>
Routine Schedule	3-dose series
Minimum Intervals	8 weeks between doses 1 & 2 6 months between doses 2 & 3
Administration	Intramuscular (IM) injection
Packaging	Vaccine packaged as 10 single-dose vials or syringes
Storage	Refrigerate between 36°F and 46°F (2°C and 8°C) - <b>Do not freeze</b>
Recommendations	<a href="#">Diphtheria, Tetanus and Pertussis (DTaP/Tdap/Td) Vaccines (ACIP) CDC's MMWR (2019)</a>
Program Letter	Not available
Billing Codes	CPT code for vaccine: 90714 CPT code for administration: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90714 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 <a href="#">Coding for Vaccine Administration (PDF)</a>
Comments	<ul style="list-style-type: none"> <li>• Several vaccines are used to prevent tetanus among children, adolescents and adults, including DTaP, Tdap, and Td.</li> <li>• Tdap can be administered regardless of interval since previous Td dose.</li> <li>• For children &lt;7 years with a contraindication to pertussis-containing vaccines, vaccine providers may administer Td for all recommended remaining DTaP doses.</li> <li>• Tdap can be used for the Td booster.</li> <li>• Td can be used as a booster every 10 years or after 5 years in the case of a dirty or major wound.</li> <li>• Td can be given at the same time as other vaccines. See <a href="#">Tetanus Prophylaxis Guide (PDF)</a>.</li> </ul>

Refer to [Vaccine Fact Sheets](#) (EZIZ.org/Resources/VaccineFactSheets).