

Topic	Penbraya™
Manufacturer	Pfizer Detailed Prescribing Information
Protects Against	Invasive meningococcal disease caused by <i>Neisseria meningitidis</i> serogroups A, B, C, W, and Y.
Routine Schedule & Intervals	<p>ACIP recommends MenACWY-MenB may be used when both MenACWY and MenB are indicated at the same visit for:</p> <ul style="list-style-type: none"> • 16 - 18 years (routine) when shared clinical decision-making favors MenB. • 10 - 25 years at increased risk: <ul style="list-style-type: none"> ○ persistent complement deficiencies, taking complement inhibitors, and anatomic or functional asplenia, including sickle cell disease. <p>Suggested routine dosing:</p> <ul style="list-style-type: none"> • dose 1 (11-12 yrs) = Menveo (MenACWY-CRM) or MenQuadfi (MenACWY-TT) • dose 2 (16-18 yrs) = Penbraya (MenACWY-TT/MenB-FHbp) • dose 3 (16-18 yrs) = Trumenba (MenB-FHbp) <p>The MenB component in Penbraya is Trumenba (MenB-FHbp). MenB brands are not interchangeable. If a patient receives Penbraya™ (MenABCWY) for MenACWY dose 2 and MenB dose 1, the MenB series must be completed with Trumenba at least 6 months later.</p> <p>Refer to: CDPH Meningococcal Vaccine Timing Guide: Routine Risk High Risk</p>
Minimum intervals	6-month interval between doses
Approved ages	10 years - 25 years of age
Administration	Intramuscular (IM) injection
Packaging	<p>Vaccine is packaged in cartons of 1, 5, and 10 kits. One kit includes one vial of Lyophilized MenACWY Component (a sterile white powder), a prefilled syringe containing the MenB Component and a vial adapter.</p> <p>To prepare vaccine for administration, reconstitute the Lyophilized MenACWY Component with the MenB Component. (Approximately 0.5mL each dose after reconstitution).</p>
Storage	<p>Before reconstitution: Refrigerate between 36°F and 46°F (2°C to 8°C). Store carton horizontally to minimize the time necessary to resuspend the MenB component. Do not freeze. Discard if the carton has been frozen.</p> <p>After reconstitution:</p> <ul style="list-style-type: none"> • Administer immediately or store between 2°C and 30°C (36°F and 86°) and use within 4 hours. Do not freeze

Vaccine Fact Sheet: MenABCWY

Topic	Penbraya™
Full ACIP Recommendations	ACIP Vaccine Recommendations
VFC Letter	VFC Penbraya Clinic Letter
Billing Codes	CPT code for vaccine: 90623 CPT code for administration: 90640 Medi-Cal Fee-For-Service (FFS) administration for VFC-supplied vaccine: 90623-SL (for ages 16 through 18 years); 90623-SL and –SK for high risk persons 10 through 15 years). ICD-10-CM code (encounter for immunization): Z23
Comments	Licensed in 2023

Refer to [Vaccine Fact Sheets](https://www.eziz.org/Resources/VaccineFactSheets) (EZIZ.org/Resources/VaccineFactSheets).