


RSV Season Immunization Recommendations

	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	April	May	Jun
Infants & Children (Clesrovimab, Nirsevimab**) 				October 1 – March 31* - All infants 0-8 months entering 1st RSV season if mother wasn't vaccinated: Clesrovimab or Nirsevimab** - High risk 8-19 month olds entering 2nd RSV season: Nirsevimab** only								
Pregnant people (Abrysvo)* 			September 1 – January 31 between 32-36 weeks gestation*									
Adults 75+ and 50-74 at increased risk (Abrysvo, Arexvy, mRESVIA) 	Offer to eligible, unvaccinated adults. CDC encourages healthcare providers to maximize the benefit of RSV vaccination by offering in late summer or early fall.											

 Recommended immunization timing

***If continuing to immunize outside recommended timeframe, make sure to:**

1. Check with insurers to ensure reimbursement. Keep remaining doses for next RSV season.
2. Unused and unexpired supply of VFC Nirsevimab, Clesrovimab or Abrysvo cannot be returned to McKesson. Label these doses as **"Keep for Fall."**

****For the 2025-2026 RSV season, IHS guidance is the preferential use of nirsevimab for eligible AI/AN infants and children.**

This publication was supported by a Centers for Disease Control and Prevention (CDC) grant. Its contents do not necessarily represent the views of CDC.

